

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Adcirca	Pulmonary Arterial Hypertension	Oral	Yes	No	Yes		No	Yes
Afinitor	Oncology- Oral/Topical	Oral	Yes	No	Yes	Protect from light; protect from moisture.	No	Yes
Alunbrig	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	No
Aranesp	Anemia	Injection SubQ or IV	Yes	No	Yes	Store at 2°C to 8°C (36°F to 46°F); do Not freeze. Do Not shake. Protect from light. Store in original carton until use. The following stability information has also been reported: May be stored at room temperature for up to 7 days	No	Yes
Avonex	Multiple Sclerosis	Injection IM	Yes	Yes	Yes	Refrigerate, Do not freeze. Protect from light. Vial: If refrigeration is not available, may be stored at 25°C (77°F) for up to 30 days	No	No
Azacitidine (Vidaza)	Oncology- Injectable	Injection SubQ or IV	Yes	Yes	Yes	Prior to reconstitution, store intact vials at room temperature of 25°C; excursions permitted to 15°C to 30°C	No	No
Betaseron	Multiple Sclerosis	Injection SubQ	Yes	Yes	No	If not used immediately following reconstitution, refrigerate solution at 2°C to 8°C and use within 3 hours; do not freeze or shake solution.	No	Yes
Bexarotene (Targretin)	Oncology- Oral/Topical	Oral	Yes	No	Yes	Store at 2°C to 25°C (36°F to 77°F). Protect from light. Avoid humidity and high temperatures after opening bottle	No	Yes
Bosulif	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	No
Bravelle	Infertility	Injection SubQ or IM	Yes	Yes	No	Lyophilized powder may be stored in the refrigerator or at room temperature of 3°C to 25°C. Protect from light. Discard unused portion.	No	Yes
Capecitabine (Xeloda)	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	Yes
Cetrotide	Infertility	Injection SubQ	Yes	Yes	No	Lyophilized powder may be stored in the refrigerator or at room temperature of 3°C to 25°C. Protect from light. Discard unused portion. Store at intact vials at 15°C to 30°C. Following reconstitution, solution is stable when refrigerated for 30 days (Novarel) or 60 days (Pregnyl).	No	Yes
Chorionic gonadotropin (Novarel, Pregnyl)	Infertility	Injection IM	Yes	Yes	No		No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Cimzia	Inflammatory Bowel Disease /	Injection SubQ	Yes	Yes	Yes	Store intact vials and syringes at 2°C to 8°C; do Not freeze. Do Not separate contents of carton prior to use. Protect from light. Bring to room temperature prior to administration. Store reconstituted solution for up to 24 hours at 2°C to 8°C (do Not freeze); Do Not leave at room temperature for more than 2 hours prior to administration.	No	Yes
Dakinza	Rheumatoid Arthritis	Oral	Yes	No	Yes	No	No	Yes
Dofetilide (Tikosyn)	Hepatitis C	Oral	Yes	No	No	None	No	Yes
Dupixent	Cardiac Disorders	Oral	Yes	Yes	Yes	N/A	No	No
	Atopic Dermatitis	Injection SubQ				Refrigerate at 2°C to 8°C. Do Not shake. Do Not freeze or store in extreme heat or cold. Store in the original carton to protect from light or physical damage. Individual autoinjectors, prefilled syringes, dose trays (containing multi-use vials and diluent syringes), or prefilled pens may be stored between 20°C and 25°C (68°F and 77°F) for a maximum single period of 14 days (Enbrel) with protection from light and sources of heat and humidity. Once an autoinjector, prefilled syringe, dose tray, or prefilled pen has been stored at room temperature, it should Not be placed back into the refrigerator; discard after 14 days (Enbrel). Dispense in original container.		
Enbrel	Psoriasis / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes		No	Yes
Epclusa	Hepatitis C	Oral	Yes	No	Yes		No	Yes
Epogen	Anemia	Injection SubQ or IV	Yes	Yes	Yes	Vials should be stored at 2°C to 8°C (36°F to 46°F). Do Not freeze. Do Not shake. Protect from light. If not used immediately following reconstitution, refrigerate solution and use within 3 hours; do not freeze or shake solution. Refrigerate. Avoid excessive or vigorous agitation; do Not shake. Discard if intact vial is left at room temperature for >12 hours prior to use.	No	Yes
Extavia	Multiple Sclerosis	Injection SubQ	Yes	Yes	No		No	Yes
Follistim AQ	Infertility	Injection SubQ	Yes	Yes	Yes		No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Forteo	Osteoporosis	Injection SubQ	Yes	Yes	Yes	Refrigerate; do Not freeze (discard if freezing occurs). Protect from light. Discard pen 28 days after first injection, even if it still contains some unused solution. Do Not use if solution is cloudy, colored, or contains solid particles.	No	Yes
Fuzeon		Injection SubQ	Yes	Yes	No		No	Yes
Ganirelix		Injection SubQ	Yes	Yes	No		No	Yes (generic)
Gattex GeNotropin Gilenya	Gastrointestinal Disorders- Other Growth Hormone & Related Disorders Multiple Sclerosis	Injection SubQ	Yes	No	Yes	Protect from light	No	No
		Injection SubQ	Yes	Yes	Yes	Refrigerated at 2°C to 8°C; do Not freeze. The carton of ancillary supplies should be stored at 25°C. After dispensing, store vials at 25°C; once dispensed, vials must be used within 90 days. Once reconstituted, store at <25°C; do Not shake or freeze; use within 3hr	No	No
		Oral	Yes	No	Yes	Protect from moisture. Dispense in original bottle; protect from high humidity and light.	No	Yes
Gilotrif	Oncology- Oral/Topical	Oral	Yes	No	Yes	Refrigerate. If needed, may store at 15°C to 30°C for up to 1 month. Avoid exposure to high temperatures; protect from intense light. Do not freeze. Discard if syringe freezes	No	No
Glatiramer acetate (Copaxone)	Multiple Sclerosis	Injection SubQ	Yes	Yes	No	Store powder refrigerated or at room temperature. Protect from light; do not freeze. Following reconstitution, multidose vials may be stored under refrigeration or at room temperature for up to 28 days.	No	Yes
Gonal-F	Infertility	Injection SubQ	Yes	Yes	Yes	Refrigerate prefilled syringes. Protect from light. Do not shake. May be removed from for a single period of up to 5 days. If not used within 5 days, the product may be returned to refrigerator up to the expiration date. Dispose of syringes if stored at room temperature for more than 5 days. Exposure to -1°C to -5°C for up to 72 hours and temperatures as low as -15°C to -25°C for up to 24 hours do not adversely affect stability. Discard unused product.	No	Yes
Granix	Neutropenia	Injection SubQ	Yes	Yes	No	Protect from light. Discard unused product.	No	Yes
Harvoni	Hepatits C	Oral	Yes	No	Yes	Protect from light. Refrigerate	No	Yes
Humantropo	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes		No	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
						Store at 2°C to 8°C (36°F to 46°F) in original container to protect from light; do Not freeze. Do Not use if frozen even if it has been thawed. Do Not store in extreme heat or cold. If needed, may be stored at room temperature up to a maximum of 25°C (77°F) for up to 14 days; discard if Not used within 14 days. Protect from moisture. Protect from light.		
Humira	Inflammatory Bowel Disease / Psoriasis	Injection SubQ	Yes	Yes	Yes	Refrigerate	No	Yes
Imatinib (Gleevec)	/ Rheumatoid Arthritis Oncology- Oral/Topical	Oral	Yes	No	Yes	Protect from moisture. Protect from light.	No	Yes
Increlex (IGF-1 Deficiency)	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Refrigerate	No	No
						Refrigerate, Protect from light. Do Not freeze or shake. After removal from the refrigerator, use within 14 days or discard.	No	Yes
Kevzara	Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes		No	Yes
						Refrigerate. Do Not freeze. Do Not shake. Protect from light. Discard any unused portion. Store in the original package	No	Yes
Kineret	Cryopyrin-Associated Periodic Syndrome / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes		No	Yes
Kisqali	Oncology- Oral/Topical	Oral	Yes	No	Yes		No	Yes
						Store under refrigeration at 2°C to 8°C. May be stored in foil pouch (opened or unopened) at room temperature of 25°C for up to 28 days. Protect from light. The colorless to pale yellow solution may darken over time if Not stored under refrigeration; however, the color change does Not affect product quality. Do Not use if solution has been stored at room temperature for >28 days.	No	No
Kitabis Pak	Cystic Fibrosis	Nebulized solution	Yes	No	No	Refrigerate. Don't freeze or shake	No	Yes
Leukine	Neutropenia	Injection SubQ or IV	Yes	Yes	No	Room temp	No	Yes
Leuprolide acetate (Lupron)	Hormonal Therapies	Injection SubQ or IM	Yes	Yes	Yes	If stored outside the original bottle, discard Orals after 30 days	No	No
Lonsurf	Oncology- Oral/Topical	Oral	Yes	No	Yes		No	No
						Orals: Store at 25°C; excursions permitted to 15°C to 30°C. Do not expose Orals to temperatures >40°C. Orals: Store at 20°C to 25°C; excursions permitted to 15°C to 30°C. Protect from moisture. Refrigerate or room temp. Protect from light. Refrigerate. Don't freeze or shake. Protect from light	No	No
Lynparza	Oncology- Oral/Topical	Oral	Yes	No	Yes		No	No
MeNopur	Infertility	Injection SubQ	Yes	Yes	No		No	Yes
Neulasta	Neutropenia	Injection SubQ	Yes	Yes	Yes		No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Neupogen	Neutropenia	Injection SubQ or IV	Yes	Yes	No	Refrigerate. Store in the original carton. Protect from light. Protect from direct sunlight. Avoid freezing. manufacturer recommends using within 24 hours due to the potential for bacterial contamination.	No	Yes
Noritropin	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Refrigerate, but do Not freeze. Protect from light. Store unopened vials in the refrigerator, away from heat and direct light. Do not freeze. Should be used within 28 days.	No	No
Nutropin AQ	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Refrigerate	No	No
Octreotide acetate (Sandostatin)	Acromegaly	Injection SubQ or IV	Yes	Yes	Yes	Refrigerate	No	Yes
Odomzo	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	No
Olysio	Hepatits C	Oral	Yes	No	Yes	Store in original bottle and protect from light. Protect from light.	No	Yes
Omnitrope	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Refrigerate	No	No
Orenica	Rheumatoid Arthritis	Injection SubQ or IV	Yes	Yes	Yes	Refrigerate, protect from light, must used within 24 hr of reconstituting	No	No
Otrexup	Psoriasis / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	No	Store intact vials and autoinjectors between 20°C and 25°C (68°F and 77°F); excursions may be permitted between 15°C and 30°C (59°F and 86°F). Protect from light. Refrigerate or room temp for 30 days. Protect from light	No	Yes
Ovidrel	Infertility	Injection SubQ	Yes	Yes	No	Refrigerate and protecting from light Intact vial: May be stored at room temperature for up to 14 days	No	Yes
Pegasys	Hepatits C	Injection SubQ	Yes	No	Yes	Prefilled syringe: May be stored at room temperature for up to 6 days. Refrigerate in the outer carton to protect from light. Time out of refrigeration should Not exceed 24 hours at 25°C. Do Not freeze. Do Not expose to extreme heat. Do Not shake.	No	Yes
Praluent	Lipid Disorders- PSK9 Inhibitors	Injection SubQ	Yes	Yes	Yes	Refrigerate in the outer carton to protect from light. Time out of refrigeration should Not exceed 24 hours at 25°C. Do Not freeze. Do Not expose to extreme heat. Do Not shake.	No	Yes
Procrit	Anemia	Injection SubQ or IV	Yes	Yes	No	Vials should be stored at 2°C to 8°C (36°F to 46°F). Do Not freeze. Do Not shake. Protect from light. Room temp. Dispense in original bottle.	No	Yes
Promacta	Immune (Idiopathic) Thrombocytopenicpurpua	Oral	Yes	No	Yes	Store at 2°C to 8°C in protective foil to protect from light. Refrigerate during transport and do Not expose to room temp for ≥24 hours.	No	Yes
Pulmozyme	Cystic Fibrosis	Nebulized Solution	Yes	No	Yes		No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Rasuvo	Psoriasis / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes	Store intact vials and autoinjectors between 20°C and 25°C (68°F and 77°F); excursions may be permitted between 15°C and 30°C (59°F and 86°F). Protect from light. Refrigerate	No	Yes
Rebetol Solution		Oral	Yes	No	No		No	No
Rebif	Multiple Sclerosis	Injection SubQ	Yes	Yes	Yes	Refrigerate; do not freeze. Protect from heat and light. Refrigeration is preferred; however, if needed, may be stored at 2°C to 25°C for up to 30 days. Refrigerate in the original carton. May also be kept at room temperature in the original carton; however, must use within 30 days (discard if Not used within 30 days). Protect from direct light and do Not expose to temp above 25°C. Do Not freeze. Do Not shake.	No	No
Repatha	Lipid Disorders- PSK9 Inhibitors	Injection SubQ	Yes	Yes	Yes	Refrigerate or room temp. Protect from light. Store at 25°C (77°F); excursions permitted between 15°C and 30°C (59°F and 86°F)	No	Yes
Repronex	Infertility	Injection SubQ or IM	Yes	Yes	No	Store at 25°C (77°F); excursions permitted between 15°C and 30°C (59°F and 86°F)	No	Yes
Ribapak	Hepatits C	Oral	Yes	No	Yes	None	No	Yes
Ribasphere	Hepatits C	Oral	Yes	No	No	None	No	Yes
Ribavirin caps (rebetol)	Hepatits C	Oral	Yes	No	Yes	None	No	Yes
Ribavirin tabs (Copegus, Moderiba)	Hepatits C	Oral	Yes	No	Yes	None	No	Yes
Rubraca	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	No
Rydapt	Oncology- Oral/Topical	Oral	Yes	No	Yes	Store in the original package to protect from moisture. Protect from light. Refrigerate	No	Yes
Saizen	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Store at 25°C; excursions permitted between 15°C and 30°C. Store at 25°C (77°F); excursions permitted to 15°C to 30°C (59°F to 86°F).	No	Yes
Samsca	Electrolyte Disorders	Oral	Yes	No	Yes	Store at 25°C (77°F); excursions permitted to 15°C to 30°C (59°F to 86°F).	No	No
Sensipar	Renal Disease	Oral	Yes	No	Yes	Refrigerate. Do not freeze. Do not shake. Store product in original carton to protect from light. Dispense in original container	No	Yes
Sildenafil (Revatio)	Pulmonary Arterial Hypertension	Oral	Yes	No	No		No	Yes
Simponi	Inflammatory Bowel Disease / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes	Dispense in original container	No	Yes
Sovaldi	Hepatits C	Oral	Yes	No	Yes	none	No	Yes
Sprycel	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Stelara	Inflammatory Bowel Disease / Rheumatoid Arthritis	Injection SubQ or IV	Yes	Yes	Yes	Refrigerate in original container to protect from light; do not freeze. Do not use if frozen even if it has been thawed. Do not store in extreme heat or cold. If needed, may be stored at room temperature up to a maximum of 25°C for up to 14 days; discard if not used within 14 days.	No	Yes
Sutent	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	Yes
Targretin GelY	Oncology- Oral/Topical	Gel	Yes	No	Yes	none	No	Yes
Tasigna	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	Yes
Technivie	Hepatitis C	Oral	Yes	No	Yes	Dispense in original container	No	Yes
Temozolomide (Temodar)	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	Yes
Tev-Tropin	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	No	Prior to reconstitution, store at 2°C to 8°C (36°F to 46°F). Following reconstitution with the provided diluent, store under refrigeration and use within 14 days (5 mg vial) or 28 days (10 mg vial); do Not freeze. Protect from light. Keep in original package.	No	No
Thalomid	Oncology- Oral/Topical	Oral	Yes	No	Yes		No	Yes
Tobramycin inhalation solution (Tobi)	Cystic Fibrosis	Nebulized Solution	Yes	No	No	Store in original package at 25°C (77°F); excursions permitted to 15°C to 30°C (59°F to 86°F). Protect from moisture. Store refrigerated. Do Not freeze or heat. After first use, store for up to 30 days at 20°C to 25°C	No	No
Tymlos	Osteoporosis	Injection SubQ	Yes	Yes	Yes		No	No
Uptravi	Pulmonary Arterial Hypertension	Oral	Yes	No	Yes		No	No
Viekira Pak	Hepatitis C	Oral	Yes	No	Yes	Dispense in original container	No	Yes
Viekira XR	Hepatitis C	Oral	Yes	No	Yes	Dispense in original container	No	Yes
Xeljanz	Rheumatoid Arthritis	Oral	Yes	No	Yes	Store at room temp	No	Yes
Xeljanz XR	Rheumatoid Arthritis	Oral	Yes	No	Yes	Store at room temp	No	Yes
Zarxio	Neutropenia	Injection SubQ or IV	Yes	Yes	No	Refrigerate in the original carton. Protect from light. Avoid freezing; if frozen, thaw in the refrigerator before administration. Discard if frozen more than once. Do not shake. Transport via a pneumatic tube has not been studied. Solutions diluted for infusion may be stored at room temperature for up to 24 hours (infusion must be completed within 24 hours of preparation). Keep in original blister pack until time of use; protect from moisture.	No	No
Zepatier	Hepatitis C	Oral	Yes	No	Yes	none	No	Yes
Zolinza	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	Yes
Zomacton	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	(see Tev-Tropin)	No	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refridgerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Zorbtive	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Store intact vials and diluent at 15°C to 30°C (59°F to 86°F). Following reconstitution with the provided diluent (bacteriostatic water for injection containing benzyl alcohol), may store under refrigeration at 2°C to 8°C (36°F to 46°F) for up to 14 days; do Not freeze. If reconstituted with sterile water for injection, solution should be used immediately.	No	No
Zytiga	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	Yes